

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

itis, 1; infantile tetanus, 1; peritonitis, 1; eclampsia, 1; hepatic cirrhosis, 2; arterio-sclerosis, 2; other causes, 3. Annual rate for week, 20.32 per 1,000.

Respectfully,

R. H. VON EZDORF,

Assistant Surgeon.

The Surgeon-General.

Report from Santiago de Cuba.

SANTIAGO DE CUBA, September 16, 1902.

SIR: Through Asst. Surg. F. E. Trotter, at Havana, I have the honor to submit the following report for the week ended September 13, 1902: Bills of health were issued to 4 vessels bound for the United States. The Norwegian steamship *Hiso* was disinfected before leaving for Mobile. No quarantinable disease has been reported. I inclose the weekly abstract of bills of health and the mortality statistics for the week.

Respectfully,

RICHARD WILSON,
Acting Assistant Surgeon.

The Surgeon-General.

[Inclosure.]

Mortality statistics for the week ended September 13, 1902, for Santiago de Cuba.

Tubercle of lungs	2
Cerebral congestion and hemorrhage	ĩ
Cerebral softening	1
Tetanus	1
TetanusOrganic diseases of the heart	3
Broncho-pneumonia	1
Diarrhea and enteritis (under 2 years)	2
Diarrhea and enteritis (2 years and over)	1
Abscess of the liver	1
Abscess of the liver	2
-	
Total	15

Annual rate of mortality for the week, 18.1 per 1,000. Estimated population, 43,000.

ECUADOR.

Quarantine measures at Guayaquil.

GUAYAQUIL, ECUADOR, August 26, 1902.

SIR: With reference to my No. 10 of the 20th instant, I have the honor to inform you that the authorities of this port have decreed that all vessels arriving from Puntarenas, Costa Rica, be admitted if provided with clean bill of health.

Vessels from Panama must submit to three days observation at Puna before granted free pratique.

That owing to there being no more cases of yellow fever at this place,

clean bills of health be issued.

The last steamship, British steamship *Peru*, from Panama, which should have arrived here yesterday, continued direct to Paita, Peru, so as to avoid the three days observation required by the port authorities. The mail will be brought here by the next steamship coming from South.

Respectfully,

TH. NAST, United States Consul.

The Assistant Secretary of State.